

ASAM/BHD PLACEMENT SUMMARY

Patient Name _____	ID # _____
Screener _____	Date ____/____/____

ASAM DIMENSION/BHD DOMAIN SEVERITY SUMMARY

No Problem (0) / Mild (1) / Moderate (2) / Significant (3) / Severe (4)

Acute Intoxication and/or Withdrawal Potential _____	Biomedical Conditions / Complications _____	Emotional, Behavioral or Cognitive Conditions and Complications _____	Readiness to Change _____	Relapse, Continued Use or Continued Problem Potential _____	Recovery Environment _____	BHD Domain: Substance Use Problems _____	BHD Domain: Independent Living Skills _____
---	--	--	------------------------------	--	-------------------------------	--	---

DIMENSION 1:

ACUTE INTOXICATION AND/OR WITHDRAWAL POTENTIAL

- Utilize CIWA-Ar score as appropriate and the ASI Alcohol / Drug Composite score
- In Substance Use Section:
 - Review individual’s previous withdrawal history
 - Review last use, amount and frequency of use and use of multiple substances in same drug class
 - Review Substance Use items related to ODs and lifetime treatment and detox

Dimension 1 Severity Rating _____

(0=no problem or stable / 1=mild / 2=moderate / 3=severe / 4=extremely severe)

- 0
- _____ Fully functioning w/ good ability to tolerate, cope with withdrawal discomfort
 - _____ No signs or symptoms of withdrawal present or are resolving; if alcohol, a CIWA-Ar of less than 3
 - _____ No signs or symptoms of intoxication
- 1
- _____ Adequate ability to tolerate or cope with withdrawal discomfort
 - _____ Mild to moderate intoxication or signs/symptoms interfere with daily functioning, but not danger to self or others
 - _____ Minimal risk of severe withdrawal resolving; if alcohol, a CIWA-Ar score of 3-7
 - _____ Sub intoxication level
- 2
- _____ Some difficulty tolerating and coping with withdrawal discomfort
 - _____ Intoxication may be severe, but responds to treatment so individual does not pose imminent danger to self or others
 - _____ Moderate signs and symptoms with moderate risk of severe withdrawal
 - _____ Somewhat intoxicated
 - _____ If alcohol, a CIWA-Ar score of 8-11
- 3
- _____ Demonstrates poor ability to tolerate and cope with withdrawal discomfort
 - _____ Severe signs and symptoms of intoxication indicating possible imminent danger to self and others
 - _____ Severe signs and symptoms or risk of severe but manageable withdrawal; or withdrawal worsening despite detoxification at less intensive level of care
 - _____ Very intoxicated
 - _____ If alcohol, a CIWA-Ar score of 12-15

- 4
- Incapacitated, with severe signs and symptoms of withdrawal
 - Severe withdrawal presents danger (e.g. seizures)
 - Continued use poses an imminent threat to life
 - Stuporous
 - If alcohol, a CIWA-Ar score of over 15

Comments (include strengths / service needs):

DIMENSION 2: BIOMEDICAL CONDITIONS / COMPLICATIONS

- Review ASI Medical Status composite score
 - In Physical Health Problems section:
 - Review Treatment for Physical Complaints
 - Review Last 30 Day and Chronic Medical Problems questions
 - Review questions regarding pregnancy status and prenatal care
 - Review Current Health Status question
-

Dimension 2 Severity Rating _____

(0=no problem or stable / 1=mild / 2=moderate / 3=significant / 4=severe)

- 0
- _____ Fully functioning with good ability to tolerate or cope with physical discomfort
 - _____ No biomedical signs or symptoms are present or biomedical problems stable
 - _____ No biomedical conditions that will interfere with treatment or create risk
 - _____ No co-occurring infections requiring treatment
 - _____ All client dental needs are being met (routine cleanings every 6 months and restorative care)
 - _____ No need for nutritional intervention
 - _____ Not pregnant
 - _____ ASI Medical Composite Score < .10
- 1
- _____ Demonstrates adequate ability to tolerate and cope with physical discomfort
 - _____ Mild to moderate signs/symptoms interfere with daily functioning but would not likely interfere with recovery treatment nor create risk
 - _____ Chronic conditions that are under control with medication and treatment (has diabetes and engaged in treatment)
 - _____ Presence of acute co-infections that are being treated
 - _____ Client is engaged in dental care at least 1 time per year and 75% of dental needs are being met
 - _____ Client has nutritional needs that are being met and client is stable or nutritional status has minimal effect on health*
 - _____ Medical condition is stable and client *is capable* of managing medical condition (refer to PPS Health Status)*
 - _____ Client is pregnant but is receiving appropriate prenatal care and not using substances
 - _____ ASI Medical Composite Score > .10
- 2
- _____ Some difficulty tolerating and coping with physical problems and/or has other biomedical problems*
 - _____ Has a biomedical problem(s) or need for medical services which may interfere with recovery treatment
 - _____ Neglects to care for serious biomedical problems*
 - _____ Acute, non-life threatening medical signs/symptoms are present
 - _____ Client non-compliant or sporadically compliant with treatment*
 - _____ Presence of acute untreated co-infections;
 - _____ Client has not been engaged in dental care for more than 1 year or has sporadic compliance with dental care plan
 - _____ Client currently engaged in complex restorative dental plan
 - _____ Client has nutritional needs that are not being addressed and health is significantly affected by nutritional status*
 - _____ Medical condition is unstable but client *is capable* of managing medical condition (refer to PPS Health Status)*
 - _____ Client is pregnant and is inconsistently receiving prenatal care, but is not using substances
 - _____ ASI Medical Composite Score > .20
- 3
- _____ Demonstrates poor ability to tolerate and cope with physical problems and/or general health is poor
 - _____ Has serious medical problems he/she neglects during outpatient treatment that require frequent medical attention (e.g., untreated diabetes)*
 - _____ Severe medical problems are present but stable
 - _____ Medical problem(s) present that would be severely exacerbated by a relapse or withdrawal
 - _____ Need 10 hrs/week or less of in home ADL assistance*
 - _____ Recent medical hospitalization (90 days)
 - _____ Client refuses dental intervention despite lack of recent engagement in care and against medical advice
 - _____ Patient or doctor report of wasting or significant observed or reported weight loss or gain in the past 3 months
 - _____ Nutritional status is profoundly affecting health*
 - _____ Client is pregnant and is receiving inconsistent prenatal care and is using substances
 - _____ Medical condition is stable and client *is not capable* of managing medical condition (refer to PPS Health Status)*
 - _____ ASI Medical Composite Score > .30
-

- 4
- _____ Incapacitated with severe medical problems*
 - _____ Severe medical problems that present life threatening risk
 - _____ Needs greater than 10 hrs/week of in home ADL assistance*
 - _____ In hospice care
 - _____ Requires assisted living*
 - _____ Emergency dental services required
 - _____ Medical condition is unstable or there are new symptoms and client *is not capable* of managing medical condition refer to PPS Health Status)*
 - _____ Client is pregnant and is not receiving prenatal care and is using substances
 - _____ ASI Medical Composite Score > .40

Comments (include strengths / service needs):

DIMENSION 3:

EMOTIONAL, BEHAVIORAL, OR COGNITIVE CONDITIONS AND COMPLICATIONS

- Review ASI Psychiatric Status composite score
 - In Mental Health Problems section:
 - Review lifetime psychiatric hospitalizations and 30 day and lifetime symptoms, particularly recent and lifetime report of hallucinations or psychotropic medication usage
 - Review Marital Status, particularly “Never Married” status
 - Review Employment Status, particularly “Retired/Disabled” status
 - Review Psychiatric Disability Status
 - Consider clinical observations during interview that client is having trouble with reality testing, thought disorders, paranoid thinking
-

Dimension 3 Severity Rating _____

(0=no problem or stable / 1=mild / 2=moderate / 3=significant / 4=severe)

- 0
- _____ No mental health problems
 - _____ No history of mental illness or use of psychotropic and addiction medications
 - _____ No lifetime psychiatric hospitalizations*
 - _____ ASI Psychiatric Composite Score of < .10
- 1
- _____ Diagnosed with prior and/or current stable mental disorder or sub threshold mental disorder (with or without prior or current use of psychotropic and addiction medications)
 - _____ Non-symptomatic or successfully treated condition*
 - _____ Mild symptoms do not impair role functioning (e.g. social, school, or work)
 - _____ Engaged in treatment and adherent or most adherent to therapy and/or psychotropic and addiction medications (include medication prescribed by general practitioner)*
 - _____ 0-1 lifetime psychiatric hospitalizations*
 - _____ ASI Psychiatric Composite Score of > .10
- 2
- _____ Emotional concerns relate to negative consequences and effects of addiction
 - _____ Suicidal ideation without plan
 - _____ Sporadically adherent to therapy and/or psychotropic and addiction medications (include medication prescribed by general practitioner)*
 - _____ Mild to moderate signs/symptoms with good response to treatment in the past; or past serious problems have long period of stability or are chronic, but do not pose high risk of harm*
 - _____ 1-4 lifetime PCS visits*
 - _____ 1-2 lifetime hospitalizations*
 - _____ ASI Psychiatric Composite Score of > .20
-

- 3
- _____ Suicidal ideation or violent impulses are potentially destabilizing but impulse control is present and impulses are not imminently dangerous, however, impulses require more than routine monitoring*
 - _____ Emotional, behavioral or cognitive problems/symptoms distract from recovery efforts and are causing moderate to severe difficulty in role functioning (e.g. school, work)*
 - _____ Frequent and/or intense symptoms with a history of significant problems that are not well stabilized but not imminently dangerous*
 - _____ History of non-adherence and currently mostly non-adherent with required psychotropic and addiction medications (include medication prescribed by general practitioner)*
 - _____ Has trouble getting along with others or has significant problems with socialization
 - _____ History of sporadic engagement in treatment or history of unsuccessful outpatient treatment*
 - _____ Uncontrolled behavior and cognitive deficits limit capacity for self-care and ADLs*
 - _____ Acute symptoms dominate clinical presentation (e.g. impaired reality testing, communication, thought processes, judgment, personal hygiene, etc.) and significantly compromise community adjustment and follow through with treatment recommendations
 - _____ Frequency of hospitalizations has increased or remained the same over the last six months and/or the length of the hospitalization has increased or remained the same*
 - _____ 4-10 lifetime PCS visits*
 - _____ 2-4 lifetime psychiatric hospitalizations*
 - _____ ASI Psychiatric Composite Score of > .30
- 4
- _____ Individual has severe and unstable psychiatric symptoms and active crisis requires secure confinement*
 - _____ Severe and acute psychiatric symptoms that pose immediate danger to self or others (e.g. imminent risk of suicide, gross neglect of self-care, psychosis with unpredictable, disorganized or violent behavior)*
 - _____ Recent history of psychiatric instability and/or escalating symptoms requiring high intensity services to prevent dangerous consequences*
 - _____ Not at all adherent with required psychotropic and addiction medications (include medication prescribed by general practitioner)*
 - _____ History of unsuccessful outpatient treatment and/or history of unsuccessful treatment at TCM level of care*
 - _____ More than 10 lifetime PCS visits*
 - _____ More than 4 lifetime psychiatric hospitalizations*
 - _____ History of detoxification visits*
 - _____ ASI Psychiatric Composite Score of > .40

Comments (include strengths / service needs)

DIMENSION 4:

READINESS TO CHANGE

- Review Major Problem, days in treatment in the last 30, and days of problems related to drugs or alcohol in the last 30 in Substance Use section
 - Review if client is awaiting charges, trial, or sentencing in Criminal Justice section
 - Review questions regarding number of minor children in out of home placement and Children’s Court in Family and Living Condition section
 - Review ASI Questions at the end of each section asking client “How serious do you feel your present... problems are?” and “How important to you now is treatment for your...?”
 - Review questions regarding recent attendance at different recovery groups
-

Dimension 4 Severity Rating _____

(0=no problem or stable / 1=mild / 2=moderate / 3=significant / 4=severe)

- 0 _____ Willingly engaged in treatment as a proactive participant, is aware of/admits to having an addiction or mental health problem and is committed to treatment for AODA use and/or mental health problems and adherence with psychiatric and addiction medications
 _____ Can articulate personal recovery goals
 _____ Willing to cut negative influences
 _____ Is in *Preparation* or *Action* Transtheoretical Stage of Change
- 1 _____ Willing to enter treatment and explore strategies for changing AODA use or dealing with mental health problems but is ambivalent about need for change (is in *Contemplation* Stage of Change)
 _____ Willing to explore the need for treatment and strategies to reduce or stop substance use or address mental health concerns
 _____ Willing to change AODA use and/or address mental health concerns but believes it will not be difficult or will not accept a full recovery treatment plan or does not recognize that he/she has an addiction or mental health problem
- 2 _____ Reluctant to agree to treatment for AODA or mental health problems but willing to be compliant to avoid negative consequences or may be legally required to engage in treatment
 _____ Able to articulate negative consequences of AODA use and/or other problematic behaviors related to mental health issues but has low commitment to change behaviors related to substance use or mental health problems
 _____ Low readiness to change and is only passively involved in treatment
 _____ Inconsistently compliant with outpatient treatment, self help or other support groups
- 3 _____ Exhibits inconsistent follow through and shows minimal awareness of AODA or mental health problems and need for treatment
 _____ Appears unaware of need to change and unwilling or only partially able to follow through with treatment recommendations
- 4 _____ Unable to follow through, has little or no awareness of AODA or mental health problems and associated negative consequences
 _____ Indifference regarding consequences of substance abuse or problematic behaviors related to mental health concerns
 _____ Not willing to explore change and is in denial regarding AODA and/or mental health problems and their implications
 _____ Disconnected from or no motivation to seek treatment against medical advice
 _____ Is not in imminent danger or unable to care for self – no immediate action required
 _____ Unable to follow through with treatment recommendations resulting in imminent danger of harm to self/others or inability to care for self – **immediate action required**

Comments (include strengths / service needs):

DIMENSION 5:

RELAPSE, CONTINUED USE OR CONTINUED PROBLEM POTENTIAL

- Review Major Problem, days in treatment in the last 30, and days of problems related to drugs or alcohol in the last 30 in Substance Use section
 - Review 30 day and lifetime medication usage in Mental Health Problems section
 - Review income source question in Education and Employment section
 - Review questions related to whether clients lives with anyone with a drug or alcohol problem and question regarding whether client has had “serious problems getting along with” in her/his support network in Family and Living Condition section
 - Review questions regarding number of minor children in out of home placement and Children’s Court in Family and Living Condition section
 - Review question regarding number of days in the past 30 of engagement in illegal activities for profit
-

Dimension 5 Severity Rating _____

(0=no problem or stable / 1=mild / 2=moderate / 3=significant / 4=severe)

- 0
- _____ No potential for further AODA or mental health problems
 - _____ Is engaged with ongoing recovery/support groups or other structured support systems
 - _____ Has positive expectancies about treatment
 - _____ Has no demographic risk factor (under 25 years of age, never married or having lived as married, unemployed, no high school diploma or GED)
 - _____ Income stable and sufficient and is able to meet monthly financial obligations
 - _____ Adequate health insurance
 - _____ No legal issues
 - _____ Internal locus of control
 - _____ No impulsivity noted
 - _____ No risk-taking or thrill-seeking
 - _____ Appropriately self-confident
 - _____ Does not or rarely requires emotional support*
- 1
- _____ One changeable demographic risk factors
 - _____ Limited gaps in service coverage
 - _____ Marginally affected by external influences
 - _____ Mostly non-impulsive
 - _____ Low level of risk-taking or thrill-seeking
 - _____ Mostly confident
 - _____ Fair self-management and relapse prevention skills
 - _____ Needs support and counseling to maintain abstinence, deal with craving, peer pressure and lifestyle and attitude changes
 - _____ Mostly adherent with prescribed psychotropic and addiction medications*
 - _____ Source of income in jeopardy frequently needs financial assistance, awaiting outcome of benefits application, or has only short term benefits*
 - _____ Occasionally requires emotional support*
-


- 2
 - Impaired recognition and understanding of substance use and/or mental health relapse issues
 - Difficulty maintaining abstinence despite engagement in treatment
 - Able to self-manage with prompting
 - One or two durable demographic risk factors
 - No income and no application for benefits, cannot cover monthly financial obligations, or bankrupt*
 - Health insurance problems (e.g., inadequate insurance, no insurance coverage, high deductibles/co-pay) resulting in limited service access and/or infrequent medical care*
 - Pending legal issues or is on parole/probation
 - Child-support issues
 - Moderately affected by external influences
 - Neither impulsive nor deliberate
 - Moderate level of risk-taking or thrill-seeking
 - Uncertain about ability to recover or ambivalent
 - Mostly adherent with prescribed psychotropic and addiction medications with failure likely to result in moderate to severe problems*
 - Regularly requires emotional support*

- 3
 - Little recognition and understanding of substance use or mental health relapse issues
 - Has poor skills to cope with addiction and/or mental health problems or to avoid or minimize relapse
 - Inconsistent or unsuccessful treatment*
 - Inconsistent and limited communication between service provider and client*
 - Three demographic risk factors
 - Immediate need for financial assistance or ineligible for benefits*
 - Substantially affected by external influences
 - Somewhat impulsive
 - High level of risk-taking or thrill-seeking
 - Dubious about ability to recover
 - Mostly non-adherent with prescribed psychotropic and addiction medications with failure likely to result in moderate to severe problems*
 - Undocumented immigrant
 - Frequency of criminal justice system involvement has increased over the last six months, or remained the same*
 - Requires frequent (more than 1 time per week) emotional support*

- 4
 - Repeated treatment episodes had little positive effect on functioning*
 - No skills to cope with addiction and/or mental health problems or to avoid or minimize relapse
 - Four or more significant demographic risks
 - Totally outer-directed
 - Very impulsive
 - Dangerous level of risk-taking or thrill-seeking
 - Very pessimistic or inappropriately confident about ability to recover

 - Not at all adherent with prescribed psychotropic and addiction medications with failure likely to result in severe problems*
 - Not in imminent danger or unable to care for self- no immediate action required
 - No skills to arrest addictive or problematic mental health behaviors or prevent relapse and continued problematic behavior places individual and/or others in imminent danger - **immediate action required***
 - Crisis involving legal system*
 - Requires daily emotional support*

Comments (include strengths / service needs):



DIMENSION 6:

RECOVERY ENVIRONMENT

- Review ASI Family/Social Status composite score
 - In the Family and Living Conditions section:
 - Review marital status items, current residence/prefers to live items, and serious conflicts with family/other people in last 30 days
 - Review items regarding living with anyone with alcohol/non-prescription drug problems
 - Review employment status in Education and Employment section
 - Review question regarding whether current living situation is supportive of recovery in Family and Living Condition section
-

Dimension 6 Severity Rating _____

(0=no problem or stable / 1=mild / 2=moderate / 3=significant / 4=severe)

- 0
- _____ Has a supportive environment or is able to cope with poor supports
 - _____ Living in a dry, drug-free home
 - _____ Stable and affordable independent housing or stable subsidized housing for greater than 1 year or has section 8 voucher*
 - _____ Few liquor outlets/no overt drug dealing
 - _____ Subcultural norms strongly discourage abusive use
 - _____ Positive leisure/recreational activities not associated with substance use or are likely to exacerbate mental health symptoms*
 - _____ No risk for emotional, physical or sexual abuse
 - _____ No logistical barriers to treatment or recovery*
 - _____ Stable relationship with children and/or dependents or does not have dependents
 - _____ Supportive significant other, friends, and family*
 - _____ Regular/periodic access to support network (church, support groups, AA, etc.)*
 - _____ Has own means of transportation consistently available*
 - _____ ASI Family/Social Composite score < .05
- 1
- _____ Has passive support in environment; family/significant other support system need to learn techniques to support recovery effort (e.g., limit setting, communication skills)*
 - _____ Significant others are not interested in supporting recovery but individual is not too distracted by this situation and is able to cope with the environment*
 - _____ Individual demonstrates motivation and willingness to obtain a positive social support system
 - _____ Safe supportive living situation in a non-dry or non drug-free home*
 - _____ Alcohol and drugs readily obtainable
 - _____ Subcultural norms discourage abusive use
 - _____ Leisure/recreational activities conducive to recovery available*
 - _____ Some risk for emotional, physical or sexual abuse
 - _____ Logistical barriers to treatment or recovery can be readily overcome*
 - _____ Occasional child care/respice needs
 - _____ Currently institutionalized; not independent but not seeking alternative*
 - _____ Transitional housing*
 - _____ Not independent but actively seeking alternative*
 - _____ Inconsistent or unreliable transportation*
 - _____ ASI Family/Social Composite score > .05
-
-

- 2
- _____ Environment is not supportive of recovery, but with clinical structure individual is able to cope most of the time*
 - _____ Living alone
 - _____ Ready access to alcohol and drugs near home
 - _____ Subcultural norms inconsistent about abusive use
 - _____ Leisure/recreational activities neutral for recovery*
 - _____ Above average risk for emotional, physical or sexual abuse
 - _____ Logistical barriers to treatment or recovery serious but resolvable*
 - _____ On-going child care/day care needs
 - _____ Not stably housed for at least 1 year*
 - _____ Grief, transition care, therapeutic intervention needed
 - _____ Single parent without support system
 - _____ Inconsistent or no dependable support system*
 - _____ ASI Family/Social Composite score > .10
- 3
- _____ Environment is not supportive of recovery and coping is difficult even with clinical structure*
 - _____ Someone in household currently dependent or abusing
 - _____ Immediate family member with substance abuse or mental health issues
 - _____ Bars/liquor stores/dealers prevalent
 - _____ Subcultural norms encourage abusive use
 - _____ Alcohol and drugs readily available at preferred leisure/recreational activities
 - _____ Substantial risk for emotional, physical or sexual abuse
 - _____ Substantial logistical impediments to treatment or recovery*
 - _____ Child abuse/neglect suspected but no evidence or involvement of Child Protective Services*
 - _____ Imminent eviction or uninhabitable home*
 - _____ Recently released from institution (last 90 days)*
 - _____ Suspected abuse by support person*
 - _____ Recent loss of primary emotional support or absent, overburdened, or poor support system*
 - _____ Lack of transportation prevents access to services*
 - _____ ASI Family/Social Composite score > .20
- 4
- _____ Environment is not supportive of recovery and is hostile and toxic to recovery or treatment progress*
 - _____ Unstable residence, living in shelter or mission, homeless*
 - _____ Subcultural norms strongly encourage abusive use
 - _____ Leisure/recreational activities pose severe risks*
 - _____ Currently being emotionally, physically or sexually abused*
 - _____ Extreme logistical impediments to treatment or recovery*
 - _____ Unable to cope with negative effects of the living environment on recovery - no immediate action required
 - _____ Environment is not supportive of recovery and is actively hostile to recovery, posing an immediate threat to safety and well-being - ***immediate action required****
 - _____ Active crisis involving dependent(s)*
 - _____ Support person is abusive*
 - _____ ASI Family/Social Composite score > .20

Comments (include strengths / service needs)

BHD DOMAIN 7: SUBSTANCE USE PROBLEMS

- Review ASI Alcohol Composite Score
 - Review ASI Drug Composite Score
 - Review ASI Psychiatric Composite Score
 - Review ASI Legal Composite Score
 - Review IV Drug Use Item on CARS Comprehensive Assessment
 - Review “Daily Activity” status, particularly with regards to “No Education, Social, or Planned Activity” status
-

BHD Domain 7 Severity Rating _____

(0=no problem or stable / 1=mild / 2=moderate / 3=significant / 4=severe)

- 0
- _____ No current or history of substance abuse issues
 - _____ No use of illicit drugs
 - _____ Greater than 1 year sobriety and actively involved in relapse prevention
 - _____ No current craving
 - _____ No other DSM 5 Substance Use Disorder criteria other than craving endorsed
 - _____ ASI ETOH Composite Score < .10
 - _____ ASI Drug Composite Score < .10
- 1
- _____ Some craving with ability to resist
 - _____ Less than 1 year sobriety
 - _____ Episodic use of alcohol (less than weekly)
 - _____ Sporadic use of drugs (less than once/week)
 - _____ Less than 2 DSM 5 Substance Use Disorder criteria endorsed
 - _____ ASI ETOH Composite Score > .10
 - _____ ASI Drug Composite Score > .10
- 2
- _____ Current use with minimal functioning impairment;
 - _____ Some craving with minimal/sporadic ability to resist
 - _____ Regular use of alcohol (once or twice a week)
 - _____ Moderate use of drugs (one to three times/week), not injected
 - _____ 2-3 DSM 5 Substance Use Disorder criteria endorsed (particularly important activities given up to use, neglect of major roles, social/interpersonal problems related to use, physical/psychological problems related to use, and time spent obtaining substances)
 - _____ Minimal detoxification history
 - _____ History of 0-1 psychiatric inpatient episodes
 - _____ History of 2-4 PCS episodes
 - _____ ASI ETOH Composite Score > .20
 - _____ ASI Drug Composite Score > .10
- 3
- _____ Current use that affects activities of daily living;
 - _____ Severe craving with minimal/sporadic ability to resist
 - _____ Active substance use causing significant functioning impairment
 - _____ Regular use of alcohol (three or more times/week)
 - _____ Frequent use of drugs (more than three times/week) and/or smoking drugs
 - _____ 4-5 DSM 5 Substance Use Disorder criteria endorsed (particularly important activities given up to use, neglect of major roles, social/interpersonal problems related to use, physical/psychological problems related to use, and time spent obtaining substances)
 - _____ History of 0-1 detoxification episodes
 - _____ History of 1-2 psychiatric inpatient episodes
 - _____ History of 4-6 PCS episodes
 - _____ ASI ETOH Composite Score > .30
 - _____ ASI Drug Composite Score >= .20
-

- 4
- _____ Severe craving with no ability to resist
 - _____ Daily intoxication
 - _____ Daily use of illicit drugs and/or IV drug use
 - _____ 6 or more DSM Substance Use Disorder criteria endorsed (particularly important activities given up to use, neglect of major roles, social/interpersonal problems related to use, physical/psychological problems related to use, and time spent obtaining substances)
 - _____ Frequency of PCS and/or Detox episodes has increased or remained the same over the last six months and/or the length of the PCS and/or Detox episodes has increased or remained the same
 - _____ History of 1-2 (or more) detoxification episodes
 - _____ History of 2 or more psychiatric inpatient episodes
 - _____ History of 6 or more PCS episodes
 - _____ ASI ETOH Composite Score > .40
 - _____ ASI Drug Composite Score >= .20

Comments (include strengths / service needs):

BHD DOMAIN 8: INDEPENDENT LIVING SKILLS

- Review medication adherence and need for medication monitoring
 - Review past year psychiatric hospitalizations
 - Review homelessness in the past year
 - Review number of arrests in the past year
 - Review degree of housing support needed
 - Review “Daily Activity” status, particularly with regards to “No Education, Social, or Planned Activity” status
 - Review Marital Status, particularly “Never Married” status
 - Review Employment Status, particularly “Retired/Disabled” status
 - Review Psychiatric Disability Status
 - Review clinical observations during interview that client is having trouble with reality testing, thought disorders, paranoid thinking or is having trouble comprehending, concentrating, remembering
-

Dimension 8 Severity Rating _____

(0=no problem or stable / 1=mild / 2=moderate / 3=significant / 4=severe)

- 0 _____ Is rated as “Independent” with regards to assistance on all the items of the Vocational and Community Living Skills sections on the MH/AODA Functional Screen*
- _____ Able to complete assistance applications and manage benefits independently*
- _____ Successfully accessing food and other benefit programs*
- _____ Does not require assistance in securing or maintaining benefits*
- _____ Regular and reliable communication between service provider*
- _____ Literate*
- _____ Can afford and is comfortable with using public or private transportation*
- _____ No cognitive impairment noted (confusion, disorientation, inattentiveness, memory problems)*
- 1 _____ Is rated as “Less than Monthly” with regards to assistance on at least 1 of the items in the Vocational and Community Living Skills sections on the MH/AODA Functional Screen*
- _____ Needs minimal or sporadic assistance to access or apply for or to maintain benefits*
- _____ Limited assistance with dependents/children required*
- _____ Requires minimal assistance in navigating system*
- _____ Low to medium level or literacy*
- _____ Client is blind/visually impaired and/or deaf/hard of hearing but can access services independently or with minimal assistance*
- _____ Has access to transportation with occasional assistance*
- _____ Cognitive impairment is absent or minimal (confusion, disorientation, inattentiveness, memory problems)*
- 2 _____ Is rated as “One to Four Times A Month” with regards to assistance on at least 1 of the items in the Vocational and Community Living Skills sections on the MH/AODA Functional Screen*
- _____ Unfamiliar with application process and/or unable to apply for or maintain benefits without significant assistance*
- _____ Requires more intensive assistance to navigate system*
- _____ Low literacy level*
- _____ Client is blind/visually impaired and/or deaf/hard of hearing and requires regular assistance to access care*
- _____ Frequently accessing transportation assistance in order to receive services (more than 3 times per month)*
- _____ Cognitive impairment is mild to moderate (confusion, disorientation, inattentiveness, memory problems)*
- _____ ASI Employment Composite score \geq .85
- 3 _____ Is rated as “More Than One Time Per Week” with regards to assistance on at least 1 of the items in the Vocational and Community Living Skills sections on the MH/AODA Functional Screen*
- _____ Illiterate*
- _____ No public or private transportation available (rural area) or client is uncomfortable using it*
- _____ Client is blind/visually impaired and/or deaf/hard of hearing and has great difficulty accessing services*
- _____ Cognitive impairment is moderate to severe (confusion, disorientation, inattentiveness, memory problems)*
- _____ ASI Employment Composite score \geq .90
-

